

# Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 01/2018

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:** PEKIRON CREAM 0.5%

**Active ingredient:** Amorolfine hydrochloride

**Dosage form:** white topical cream

**Print on wrapping:** ペキロンクリーム 0.5% 10g, PEKIRON CREAM



## Effects of this medicine

This medicine selectively disturbs two-stage synthetic pathway of fungal cell membrane to damage the structure and function of the cell membrane, and shows antibacterial action.

It is usually used to treat skin fungal infection such as ringworm (tinea, athlete's foot etc.), candidiasis of skin and chromophytosis.

## Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, apply on the affected area once a day. Strictly follow the instructions.
- Be careful not to put this medicine in eyes, and do not use it on cornea and conjunctiva.
- If you miss a dose, apply as soon as possible.
- If you accidentally use more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop using this medicine unless your doctor instructs you to do so.

## Precautions while taking this medicine

## Possible adverse reactions to this medicine

The most commonly reported adverse reactions include contact dermatitis, redness, itch and erythema. If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

No pertinent entries.

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

## Storage conditions and other information

- Keep out the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

**For healthcare professional use only** / /

For further information, talk to your doctor or pharmacist.